

Micford Meadow Centre ATAA 240217  
24/02/17

Samples used = 10

Species name	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	Sum	SCVI	BI
GPS easting / ST	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
GPS northing / ST	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
0: Brachythecium rutabulum	1	1	1	1	1	1	1	1	1	1	1										10	2	80
1: Kindbergia praelonga	1	1	1	1	1	1	1	1	1												7	2	42
2: Barbula convoluta					1	1	1	1													2	3	8
3: Oxynchium hians					1	1	1	1	1												4	3	28
4: Phascum cuspidatum					1	1															2	4	0
5: Bryum capillare							1														1	2	5
6: Dicranoweisia cirrata							1														1	3	3
7: Frullania dilatata							1														1	3	3
8: Hypnum resupinatum							1														1	3	0
Summary	2	2	2	2	1	5	4	7	2	3	1	0	0	0	0	0	0	0	0	0	29	2.778	169

Species Richness = 9  
Shannon-Wiener Index = 1.8167(1.3895)  
Simpson Index = 4.7514(3.1541)  
Berger-Parker Dominance Index = 0.3448(0.4733)  
Density = 0.029 per sq.m.  
Species Conservation Value Index = 2.7777+/-0.6285  
Biomass Index = 169.0

Chao1 (pop.) Richness = 13.0+/-4.2573  
Chao2 (pres./abs.) Richness = 13.0+/-4.2573  
Bootstrap Richness = 11.4261+/-2.7738  
Jackknife Richness = 12.6+/-11.3841

